



HIRA-mediated H3.3 deposition is required for de novo paternal nuclear envelope formation in mouse zygotes

Citation

Inoue, Azusa, and Yi Zhang. 2013. HIRA-mediated H3.3 deposition is required for de novo paternal nuclear envelope formation in mouse zygotes. *Epigenetics & Chromatin* 6(Suppl 1): 05.

Published Version

doi:10.1186/1756-8935-6-S1-05

Permanent link

<http://nrs.harvard.edu/urn-3:HUL.InstRepos:10609732>

Terms of Use

This article was downloaded from Harvard University's DASH repository, and is made available under the terms and conditions applicable to Other Posted Material, as set forth at <http://nrs.harvard.edu/urn-3:HUL.InstRepos:dash.current.terms-of-use#LAA>

Share Your Story

The Harvard community has made this article openly available.
Please share how this access benefits you. [Submit a story](#).

[Accessibility](#)

ORAL PRESENTATION

Open Access

HIRA-mediated H3.3 deposition is required for de novo paternal nuclear envelope formation in mouse zygotes

Azusa Inoue^{1,2,3*}, Yi Zhang^{1,2,3}

From Epigenetics and Chromatin: Interactions and processes
Boston, MA, USA. 11-13 March 2013

Paternal genome undergoes dynamic chromatin remodeling after fertilization. The first event is the exchange of protamines with maternally deposited histones including H3.3. Although HIRA has been shown to be responsible for H3.3 deposition in invertebrates, whether such function is conserved in vertebrates is not known. Furthermore, the biological significance of the protamine-histone exchange is also unknown. To address these questions, we depleted the maternal pool of HIRA protein by using an *in vitro* oocyte growth system where siRNA is injected into small growing oocytes with surrounding granulosa cells. In HIRA-depleted zygotes, H3.3 is not incorporated into the paternal genome while protamines are removed, indicating that protamine-removal is uncoupled from H3.3 deposition. Inhibition of H3.3 deposition prevents the de novo nucleosome assembly and the formation of paternal pronucleus. Immunostaining of nuclear envelope components revealed the lack of Lamin B1 and nuclear pore complex assembly and the defect in flattening of endoplasmic reticulum (ER) sheets, the precursor of nuclear envelope, only in the paternal genome of HIRA-depleted zygotes. Furthermore, the HIRA-depletion prevented the localization of Ran, a GTPase required for ER fusion, in the paternal genome. Together, our results not only provide the first direct evidence that HIRA is responsible for H3.3 deposition of the paternal genome in vertebrates, but also suggest that HIRA-mediated nucleosome assembly is a pre-requirement for the recruitment of factors, such as Ran GTPase, involved in ER flattening and nuclear envelope formation during de novo paternal pronucleus formation.

Author details

¹Howard Hughes Medical Institute, USA. ²Program in Cellular and Molecular Medicine, Boston Children's Hospital, USA. ³Department of Genetics & Pediatrics, Harvard Medical School, 200 Longwood Avenue, Boston, MA 02115, USA.

Published: 18 March 2013

doi:10.1186/1756-8935-6-S1-O5

Cite this article as: Inoue and Zhang: HIRA-mediated H3.3 deposition is required for de novo paternal nuclear envelope formation in mouse zygotes. *Epigenetics & Chromatin* 2013 **6**(Suppl 1):O5.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Howard Hughes Medical Institute, USA
Full list of author information is available at the end of the article